

Implementing a multi-modal support service model for family caregivers looking after persons with age-related macular degeneration: study protocol for a randomised control trial.

Bamini Gopinath^{1,2}

Ashley Craig²

Annette Kifley^{1,2}

Gerald Liew¹

Jaye Bloffwitch¹

Kim Van Vu^{1,2}

Nichole Joachim¹

Rob Cummins⁴

Julie Heraghty⁴

Timothy Broady⁵

Alison Hayes³

Paul Mitchell¹

¹Centre for Vision Research, Department of Ophthalmology and Westmead Institute for Medical Research, University of Sydney, NSW, Australia.

²John Walsh Centre for Rehabilitation Research, Sydney Medical School, Kolling Medical Research Institute, University of Sydney, Sydney, Australia.

³Sydney School of Public Health, University of Sydney, Sydney, Australia.

⁴Macular Disease Foundation Australia, Sydney, Australia.

⁵Carers NSW, Sydney, Australia.

Correspondence to:

Associate Professor Bamini Gopinath

Centre for Vision Research

Westmead Hospital, Westmead, NSW 2145, Australia.

Telephone: 61-2-8627 3368 Fax: 61-2-8627 3099

Email: bamini.gopinath@sydney.edu.au

Abstract: 297; Text: 3,714; Tables: 2; Figure: 1

ABSTRACT

Introduction: Age-related macular degeneration (AMD) is a leading cause of blindness and low vision among older adults. Previous research shows a high prevalence of distress and disruption to lifestyle among family caregivers of persons with late AMD. This supports existing evidence that caregivers are ‘hidden patients’ at risk of poor outcomes. There is ample scope for improving the support available to caregivers, and further research should be undertaken into developing services that are tailored to the requirements of family caregivers of persons with AMD. This study aims to implement and evaluate an innovative, multi-modal support service program that aims to empower family caregivers by improving their coping strategies, and enhancing feelings of self-efficacy, and helping them make the most of available sources of social and financial support.

Methods and analysis: A randomised controlled trial consisting of 360 caregiver-patient pairs (180 each in the intervention group and wait-list control group). The intervention group will receive the following: 1) mail-delivered cognitive behavioural therapy (M-CBT), designed to improve psychological adjustment and adaptive coping skills; 2) telephone-delivered group counselling sessions allowing carers to explore the impacts of caring and share their experiences; and 3) education on available community services/resources, financial benefits, and respite services. The CBT embedded in this program is the best evaluated and widely used psychosocial intervention. Primary outcome is reduction in caregiver burden. Secondary outcomes include improvements in caregiver mental wellbeing quality of life, fatigue and self-efficacy. Economic analysis will inform whether this intervention is good value for money, and if it is feasible to roll out this service on a larger scale.

Ethics and dissemination: The study was approved by the University of Sydney Human Research Ethics Committee (HREC). Study findings will be disseminated via presentations at national/international conferences, and peer-reviewed journal articles.

Trial registration number: ACTRN12616001461482

Strengths and limitations

- It is an innovative study as it is the first to implement and evaluate a comprehensive support service tailored to family caregivers of persons with AMD.
- Randomised controlled trial design with a wait-list control group.
- Translation of evidence-based interventions to practice - there is evidence that CBT can be effective in reducing caregiver burden and distress; and it is the best evaluated and widely used psychosocial intervention.
- It is a multi-component intervention and prior research has shown these are more effective than single interventions, because it uses several techniques and address a variety of caregiver needs.
- Recruitment of participants was confined to only one state in Australia.
- Blinding of the intervention is not possible to participants and project staff (i.e. study coordinator).

INTRODUCTION

Age-related macular degeneration (AMD) is a progressive, chronic disease of the central retina, and is a leading cause of blindness and low vision among older adults.¹ Advanced age-related macular degeneration, including neovascular age-related macular degeneration (wet) and geographic atrophy (late dry), is associated with substantial, progressive visual impairment.² Visual impairment from AMD often means that patients have reduced ability to engage in everyday activities and family members are often called upon to provide physical and emotional support.³⁻⁶ Caring for loved ones with vision loss is burdensome and often leaves the carer exhausted and at risk of health problems.^{5,7,8} A UK study⁹ showed an elevated level of caregiver burden associated with caring for family with wet AMD, which was equivalent to the burden experienced by caregivers of rheumatoid arthritis and multiple sclerosis patients,^{10,11} and higher than for colorectal cancer patients.⁹

We had previously conducted a cross-sectional survey of 500 caregivers of persons with wet AMD, and showed a higher than expected prevalence of emotional distress (>50%; n=280), including feeling sad, frustrated and/or isolated in this cohort.⁷ This finding underscored the difficulty of coping with the challenges related to assisting persons with advanced AMD. If undetected and untreated, negative outcomes related to caregiver distress could persist or worsen over time.⁵ Further, healthcare providers are often forced to neglect caregivers and place priority on service delivery for their patients based on budgetary and strategic planning priorities.¹² This could lead to increased anxiety/ stress, resulting in greater overall burden in informal carers.¹²

Over half of surveyed caregivers of people with wet AMD in our study (n=295) also neglected their own needs, and personal or family interests and had to make changes to one or more of areas of their life (e.g. retirement plans).⁷ This self-perceived disruption to the social support they receive from their informal network of family and friends, and the formal

network (e.g. paid employment), as a result of caring for someone with AMD could perpetuate negative outcomes in caregivers.⁷ The qualitative analyses component of our study,¹³ showed that significant sacrifices were made by the caregiver in order to meet the needs of the care recipient with late AMD, which could also add to the burden and distress experienced by the carer. Finally, our study showed that very few family caregivers sought or received respite, and this added burden can have a negative impact upon the relationship between caregiver and care recipient.¹³

Despite the importance of information and support to help family caregivers; interventions to increase support for family caregivers have lagged far behind those provided for patients.¹⁴ There is ample scope for improving the support available to caregivers, and further research should be undertaken into developing services that are tailored to the requirements of family caregivers.¹⁵ Early intervention, awareness programs and coordination of community resources could alleviate carer distress.⁷ There is strong consensus that once information needs are met, family caregivers are likely to benefit from additional interventions such as improving their problem solving skills and extending their social support network.¹⁶ Specifically, psychosocial interventions such as cognitive behavioural therapy (CBT) can decrease caregiver distress, by fostering a more efficient, systematic approach to caregiving that enhances motivation, and self-efficacy when providing care for a family member.^{5,17-22} CBT is a recognised efficacious strategy, and it aims to help the client to develop skills that they can use outside of the therapy session and continue using when therapy ends.¹⁷⁻²⁰ Moreover, mail-delivered CBT (M-CBT) involving written materials being posted to older caregivers is optimal, as it allows them to review the materials as often as needed and at their own pace.²³ In randomised clinical trials (RCTs), M-CBT has established efficacy against control in a wide range of disorders.^{24,25}

The proposed intervention study is novel as it will be the first to implement and evaluate a comprehensive support service tailored to family caregivers of individuals with AMD. The intervention includes M-CBT and telephone-delivered group therapeutic program (facilitated by a professional counsellor), which will effectively consolidate the psychoeducation, as well as educational materials on community support, respite services and carer financial entitlements and benefits. The study will be a two-arm RCT trial with an economic evaluation. This study was co-developed with key stakeholders or partner organisations – Macular Disease Foundation of Australia (MDFA) and Carers NSW. MDFA is the national peak body representing people with macular disease, their families and carers; and Carers NSW is a state-wide organisation representing informal carers. The current study aims to improve the design and delivery of these organisation's existing support services and programs, in a way that makes it easier for family carers to have timely access to a coordinated, multi-component intervention targeting drivers of caregiver stress and burden. A unique aspect of this study is that MDFA staff will be trained in the principles of CBT. MDFA currently run support services from their Sydney (Australia) office for clients with AMD and their family members and/or carers; and this intervention will be implemented as part of these existing support services. There is evidence to support lay providers without any healthcare experience delivering CBT, under the supervision of experts.²³ Appropriate training of MDFA staff will ensure that CBT becomes a relatively stable and enduring part of the behavioural repertoire of MDFA in the future.

The primary hypotheses assert that the multi-modal intervention will lead to significant improvements in caregiver burden (primary study outcome), health related quality of life, mental and emotional wellbeing, fatigue, self-efficacy and life satisfaction (secondary outcomes). A high rate of adherence and participant satisfaction with the multi-modal intervention is also hypothesised, with delivery both acceptable and feasible to stakeholder

organisations. Finally, the intervention is hypothesised to be cost-effective when compared with usual care, when considering health outcomes for family caregivers.

METHODS

Study design and participants

This ongoing study is designed as a multisite, two-arm randomised controlled trial (RCT); with an intervention group and a waitlist control group (Fig 1). The protocol is developed in accordance with the Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) recommendation.²⁶ It is a 3-year study, and recruitment commenced on January 2017. This study aims to recruit participant pairs comprising a family caregiver (related to an individual diagnosed with AMD) and the care recipient (individual diagnosed with AMD). A family caregiver will be defined as “any person who, without being a professional or belonging to a social support network, and in some way, is directly implicated in the patient’s eye care or is directly affected by the patient’s health problem”.⁹

Eligibility of participants

Inclusion criteria

1. Adult aged >18 years.
2. Family caregiver to an individual diagnosed with AMD and related to the care recipient (i.e., spouse, child, or sibling).
3. Willing to engage in a 10-week therapeutic intervention over a 3-month period.

Exclusion criteria

1. Family caregiver lacking sufficient fluency in spoken English to engage in therapy.

Recruitment

Participant pairs will be recruited primarily from private eyes clinic in Westmead and Liverpool (Sydney, New South Wales, Australia). Recruitment involves an opt-in two-step process for a participant pair. Consent for participation is required from both the caregiver and care recipient. Step 1 involves screening for interest in participating in the research via either: (1) the study co-ordinator discussing the study with potential participants attending the eye clinic; or (2) advertising the study through MDFA and Carers NSW newsletters, and clients self-referring as a result of the provision of information. The advertisement will identify “possible emotional strain and stress” associated with caring, and the treatment’s aims of “improving emotional well-being and coping”.

Step 2 involves providing further information about the study via the study information sheet and obtaining written consent for the participant by either: (1) Providing the information sheets, consent forms and baseline questionnaires to potential participant pairs attending the eye clinic – this will be completed at the clinic or mail-returned with a reply-paid envelope; or (2) Telephoning the potential participant to discuss the research further, followed by posting information sheets, consent forms and baseline questionnaires to people who have indicated willingness to participate. Loss of contact will be minimised by asking for multiple contact methods and sending a letter at regular intervals.

Randomisation and blinding

All participants will be randomly assigned to one of two groups: (i) multi-modal treatment or (ii) wait-list control (Fig 1). The randomisation sequence will be generated centrally using permuted blocks of mixed size, stratified by recruitment source (clinic versus self-referral as a result of responding to MDFA and Carers NSW advertisements), to ensure equal numbers while maintaining an unpredictable sequence. Assignments to the intervention or control group will be managed centrally, separately from recruiting and treating staff, in order to

ensure that a randomisation number and group is assigned and recorded for each recruit before their allocation becomes apparent.

Intervention group

The intervention group will receive a multi-modal support service program, consisting of: (1) brief M-CBT treatment to be delivered fortnightly as five individual modules; and (2) five Talk-Link group counselling sessions (an existing Carers NSW Support Service). The Talk-Link Counselling and M-CBT will occur weekly on an alternating basis (i.e., week one: M-CBT, week two: Talk Link, week 3: M-CBT, and so on). The whole intervention will be conducted over 10 weeks (Table 1).

The M-CBT of fortnightly modules (Table 1) formatted in Microsoft PowerPoint, with additional homework worksheets and accompanying templates for practicing acquired skills will be mailed to participants in the intervention group. Each module will target a specific stressor and/or train a new adaptive coping method (Table 1); and will be supported by targeted homework assignments for the caregiver to practice between sessions. These will draw upon standard CBT practice and on a number of additional sources including published trials, empirical studies, and treatment guides.¹⁷⁻²⁰ Participants will be telephoned following each M-CBT module for a brief conversation (5-10 mins) aimed at: (i) addressing any queries regarding the information presented in the most recent module; (ii) encouraging the learning and practicing of skills from the previous module; (iii) normalising the challenges of acquiring new skills; (iv) emphasising the importance of consistent practice of newly acquired skills for positive impact; and (v) reminding participants in the case of unread materials. We will also ask participants how much time they spent on each module that week including reading the materials and completing the related homework sheets. Calls will be logged for each participant. The educational materials sent out to participants will aim to

increase their knowledge about support and respite services, and financial entitlements and benefits, how best to access them, which has been previously shown to increase carer sense of competence and reduce depression.¹⁴

Participants in the intervention group will be asked to complete a form prior to participating in the Carers NSW Talk-Link group sessions to identify the issues they might want to discuss (Table 1). These will then be developed into weekly topics to guide group discussions. Information/ newsletters will be provided by the facilitators. Using teleconferencing, a group of 6-8 caregivers and two trained facilitators will get together over the telephone, at the same time each week, to explore issues around caring for someone with AMD. These sessions will last for 1 hour.

Control group

During the treatment phase, the active waitlist control group will receive reading materials concerning AMD and caring for persons diagnosed with the condition. All control participants will be offered the opportunity to receive the multi-component intervention after the study ends (12-18 months after inclusion).

Measurement of covariates

A pre-intervention baseline questionnaire will obtain demographic and health related information (i.e. covariate data) from all family caregivers: age, sex, years of education, employment status, living arrangements, marital status, whether they are the sole caregiver, relationship to care recipient, and their own health status (e.g. subjective self-rated health and any chronic conditions that they suffer). Follow-up assessments will be conducted by postal questionnaires for all participants (caregivers only and not care recipients) from both RCT

arms: immediately after completion of the 10-week program, and 6 and 12 months after initiation of the intervention (Fig 1).

General caregiving activities will be assessed by asking the caregiver whether they performed any of the basic (e.g. toileting, feeding, dressing, grooming) and instrumental daily living activities (e.g. housekeeping, food preparation, shopping, telephone usage, and transportation) for their care recipient using a validated questionnaire.²⁷ In relation to each of these activities they will be asked to report whether they gave: (a) No help or little help given; (b) Moderate amount of help given; (c) High amount of help given; or (d) Not applicable. For eye-related activities that the caregiver assisted with, questions will quantify the impact of the anti-VEGF therapy received by the care recipient: the number of appointments attended; the average time taken for appointments; time taken away from work; loss of income; and whether they administered eye drops to the patient post-injection. Caregivers will be asked to detail the frequency (1 day/week; 2-3 days/week; 4-6 days/week; 7 days/week; 1-3 days/month or other) and duration of care (<1 hour; 1-2 hours; 3-5 hours; 5-8 hours; >8 hours or other) that they generally provide. The caregiver will also report the level of dependency of the care recipient on them, by choosing one of the following: not at all dependent; somewhat dependent; moderately dependent; very dependent; or extremely dependent.

The following information related to care recipients (patients with AMD) will be collected through the eye clinic records: visual acuity in better eye, number of anti-vascular endothelial growth factor (VEGF) injections to date, number of clinic appointments and total follow-up period. A brief questionnaire will also be administered to care recipients at baseline, in order to collect information on demographic details (age, sex, ethnicity, education, employment status, living arrangements, and marital status); any other eye diseases (e.g. cataract, glaucoma) and chronic conditions they may suffer from e.g. heart

disease, stroke, diabetes, arthritis. The questionnaire will also determine self-reported vision-related health status, using the National Eye Institute Visual Function Questionnaire (NEI-VFQ-25).²⁸ This is a 25-item scale used to generate a single composite score from 0 to 100 (100 is maximum visual function).⁹

Outcome measurement

Outcome data will be obtained from all randomised study participants using validated questionnaires or tools at baseline, 10 weeks, 6 and 12 months. Subjective caregiver burden is the primary outcome and will be measured using the Caregiver Burden Scale, CBS (Table 2). Secondary outcomes i.e. fatigue, depressive symptoms, general health (EQ-5D-5L), and self-efficacy are also detailed in Table 2. Further, caregivers will indicate their overall quality of life on a linear analog uniscale from 0 (worst possible) to 10 (best possible). Caregivers will be asked to identify whether caring for a family member with AMD created any specific problems, including: feeling tired, anxious, stressed, sad and/or depressed; impediments to employment, retirement, social activities and travel plans; and/or relationship problems as a result of the caregiving experience.

Individual participant consent will be sought from caregivers for Medicare benefits scheme (MBS) data linkage. Permission will be sought to link data from 12 months prior to the intervention and up until one year after the intervention. Hence, health-related resource use data e.g. medical consultations and prescription use will be compared between the 2 RCT arms and will contribute to the economic evaluation. To protect the participants' privacy, all outcome data will be kept in a separate password-protected file from the document containing the participants' names and identification codes. All documents related to the study will be stored in restricted-access facilities or locked cupboards, and on restricted-access servers and password-protected PC's and files as per University of Sydney HREC requirements.

Acceptability and feasibility of the intervention

Treatment acceptability in the intervention arm will be assessed immediately after completion of the 10-week multi-modal treatment by asking participants to rate their satisfaction with the: (1) overall support service program they received; (2) CBT modules they completed; and (3) adherence to the program. Participants will respond to these 3 questions using a 5-point Likert scale, ranging from 'Very Satisfied' to 'Very Dissatisfied'. Participants will also be asked to respond as yes/no to the following questions: (1) 'Would you feel confident in recommending this treatment to a friend?' (2) 'Was it worth your time participating in this program?' Finally, participants will be asked to specify which components of the program they found most useful or valuable; and suggestions will be elicited on how the program could be improved further. Moreover, the percentage of people who enter the study compared to the total number to whom participation is offered, and the follow-up rate achieved will also be determined. Treatment feasibility will be broadly assessed based on preliminary outcomes achieved; cost-effectiveness of the intervention, combined with the ease with which support staff could support participants.

Economic evaluation

The analysis will be conducted from societal perspective and will include costs incurred by individuals and healthcare providers. This perspective will be taken because the cost consequences of this intervention may extend beyond the domain of healthcare to deliver the model and the cost-effectiveness compared. This will be a 'within-trial' economic evaluation carried out within the timeframe and context of the trial, and will determine the cost-utility of this support service compared with usual care. We will determine the cost per person to deliver the support service using a bottom up approach. This will include the cost of training MDFA staff, telephone consults, educational materials and other consumables. These will be

valued at market rates and presented in 2016 \$A. This will enable us to determine the cost of wider roll out of the program. Direct health care costs will be determined through patient-level linkage to MBS and the PBS. Healthcare resource use during the trial will include (1) primary and secondary health care consultations and admissions; (2) use of medications, including anti-depressants and anxiolytic medications. Healthcare costs will include both government contributions to healthcare plus any patient out-of-pocket expenses. We will also measure and value time contributions of caregivers to attend the course and take part in discussion groups, using median wage rates.

Participants' health states will be captured using the 5 domains of the EQ-5D-5L namely, mobility, self-care, usual activities, pain/discomfort and anxiety/depression. The responses on each domain of the EQ-5D-5L will be converted into utility values using a valuation algorithm for the Australian population.²⁹ Quality adjusted life years (QALYs) will be determined over the duration of the trial from the summation of mean utility x time interval between EQ-5D administration times. The incremental cost-effectiveness ratio in AUD/ QALY will be determined in the intervention compared with control. Bootstrapping will be used to estimate a distribution around mean costs and outcomes, and to calculate the confidence intervals around the incremental cost effectiveness ratio. One-way sensitivity analysis will be conducted around key variables, and a probabilistic sensitivity analysis will be conducted to estimate the joint uncertainty in all parameters. A cost-effectiveness acceptability curve will be derived which summarizes the probability of the intervention being cost-effective based on willingness to pay per QALY gained.

Sample size considerations

We plan to recruit 360 caregiver-patient pairs (180 in each RCT arm) over 4-6 months.

Allowing 20% loss to follow up due to changes in the status of care recipients or death/

dropout of carers, we will have 288 carers available for analysis. For our primary endpoint (caregiver burden scores), 194 available caregivers provides 80% power to detect a moderate and clinically meaningful effect size of 0.5 of the intervention on caregiver burden, after allowing for full compliance of 81%, at $\alpha=0.05$. An 81% full compliance level represents 90% of recruits in the intervention arm actively taking up the educational and support program, among whom 90% actively practice the strategies learned. We have chosen to recruit the higher number of 288 available caregivers, so that we will additionally be able to estimate rates of uptake, acceptance, satisfaction and improvement with intervention with high precision (standard errors <6%), and have well-powered secondary analyses including an effect size of 0.5 of intervention on mean overall QOL, and a difference of 20% vs 40% between intervention and control groups in the proportion reporting specific negative impacts of caregiving.

Data collection and analysis plan

All data will be collected using questionnaire booklets and data collection sheets, and will be subsequently entered into a secure online platform, called Research Electronic Data Capture (REDCap). Primary and secondary study endpoints will be analysed by intention-to-treat.

The primary study endpoint will be post-intervention change from baseline in CBS scores.

The primary analysis will test for difference between intervention and control groups in post-intervention change from baseline using linear mixed models for repeated measures.

Secondary analyses will directly test change in scores at each follow-up within the intervention and control groups; and test for difference between intervention and control groups in the average of the three post-intervention CBS scores. The percentage of individuals within intervention and control arms who demonstrate a large change in caregiver burden, defined as a change of half a standard deviation or more, will be estimated with 95%

confidence intervals (CI). Secondary endpoints are: percentage of participants reporting overall satisfaction with the intervention and each major subcomponent, percentage of participants completing the overall program and each major subcomponent, change from baseline in quality of life, fatigue and self-efficacy scores, and post-intervention frequency of depressive symptoms.

Differences between the intervention and control groups in individual EQ-5D-5L domain and depression outcomes will be tested using χ^2 tests, and mean post-intervention level of caregiver burden, fatigue, self-efficacy, depressive symptoms, and general health will be tested using two-sample *t*-tests. Rates of uptake, continued participation, acceptability and satisfaction with intervention will be estimated with 95% CIs. Associations between baseline recipient and caregiver characteristics, treatment experience and post-intervention outcomes will be explored using linear mixed models for repeated measures. Baseline characteristics of the intervention and control groups will be examined.

Safety measures and endpoint

A clinical psychologist (AC), experienced in the assessment and treatment of mental health disorders is supporting this study. The occurrence of adverse events, such as clinical psychological presentation (i.e., anxiety, depression), or risk of self-harm and suicide, are considered safety end-point measures of this trial. Regular contact by support staff will identify whether the safety endpoint measure is required for any participant and the clinical psychologist on the research team, as per accepted mental health clinical guidelines, would instigate appropriate action.

ETHICS AND DISSEMINATION

The study was approved by the University of Sydney Human Research Ethics Committee. The committee deemed this study as low risk in terms of ethical issues. The written papers from this study will be submitted for publication in quality peer-reviewed medical and health journals. Study findings will also be disseminated via presentations at local, national and international conferences.

Contributors: The authors' responsibilities were as follows—BG, AC, AK, GL, AH, PM, RC, JH and TB: study concept and design; BG, JB, KVV and NJ: acquisition of data; JB, BG: drafting of the manuscript; BG, AC, AK, GL, AH, PM, JB, KVV, NJ, RC, JH and TB: critical revision of the manuscript. All authors have given final approval of the version to be published.

Funding: The study is supported by an Australian National Health and Medical Research Council Partnership Project Grant (APP1115729).

Competing interests: None declared.

Ethics and dissemination: The University of Sydney Human Research Ethics Committee approved the study (reference: HREC 2016/793). Findings will be disseminated through research conferences and peer-reviewed journals.

Data sharing statement: Study data will be available on request to BG once the research team has completed the pre-planned analyses.

REFERENCES

1. Foran S, Wang JJ, Mitchell P. Causes of visual impairment in two older population cross-sections: the Blue Mountains Eye Study. *Ophthalmic Epidemiol.* 2003;10:215-225.
2. Lim LS, Mitchell P, Seddon JM, Holz FG, Wong TY. Age-related macular degeneration. *Lancet.* 2012;379:1728-1738.

3. Hassell JB, Lamoureux EL, Keeffe JE. Impact of age related macular degeneration on quality of life. *Br J Ophthalmol*. 2006;90:593-596.
4. Williams RA, Brody BL, Thomas RG, Kaplan RM, Brown SI. The psychosocial impact of macular degeneration. *Arch Ophthalmol*. 1998;116:514-520.
5. Bambara JK, Owsley C, Wadley V et al. Family caregiver social problem-solving abilities and adjustment to caring for a relative with vision loss. *Invest Ophthalmol Vis Sci*. 2009;50:1585-1592.
6. Gopinath B, Liew G, Burlutsky G, Mitchell P. Age-related macular degeneration and 5-year incidence of impaired activities of daily living. *Maturitas*. 2014;77:263-266.
7. Gopinath B, Kifley A, Cummins R, Heraghty J, Mitchell P. Predictors of psychological distress in caregivers of older persons with wet age-related macular degeneration. *Aging Ment Health*. 2014;1-8.
8. Hu C, Kung S, Rummans TA, Clark MM, Lapid MI. Reducing caregiver stress with internet-based interventions: a systematic review of open-label and randomized controlled trials. *J Am Med Inform Assoc*. 2014.
9. Gohil R, Crosby-Nwaobi R, Forbes A et al. Caregiver Burden in Patients Receiving Ranibizumab Therapy for Neovascular Age Related Macular Degeneration. *PLoS ONE*. 2015;10:e0129361.
10. Orgul S, Gaspar AZ, Hendrickson P, Flammer J. Comparison of the severity of normal-tension glaucoma in men and women. *Ophthalmologica*. 1994;208:142-144.
11. Varma R, Tielsch JM, Quigley HA et al. Race-, age-, gender-, and refractive error-related differences in the normal optic disc. *Arch Ophthalmol*. 1994;112:1068-1076.
12. Westaway L, Wittich W, Overbury O. Depression and Burden in Spouses of Individuals with Sensory Impairment. *Insight: Research and Practice in Visual Impairment and Blindness*. 2011;4:29-36.

13. Vukicevic M, Heraghty J, Cummins R, Gopinath B, Mitchell P. Caregiver perceptions about the impact of caring for patients with wet age-related macular degeneration. *Eye (Lond)*. 2015.
14. Reinhard SC, Given B, Petlick NH, Bemis A. Supporting Family Caregivers in Providing Care. In: Hughes RG, ed. *Patient Safety and Quality: An Evidence-Based Handbook for Nurses*. Rockville (MD): Agency for Healthcare Research and Quality (US) 2008:341-404.
15. Deloitte Access Economics. *The economic value of informal care in Australia in 2015*. 2015. Deloitte Access Economics Pty. Ltd. 2015.
16. Williams K, Owen A. A contribution to research and development in the carer support sector: Lessons on effective caring. *Family Matters*. 2009;82:38-46.
17. Craig A, Hancock K, Chang E, Dickson H. The effectiveness of group psychological intervention in enhancing perceptions of control following spinal cord injury. *Aust N Z J Psychiatry*. 1998;32:112-118.
18. Horrell L, Goldsmith KA, Tylee AT et al. One-day cognitive-behavioural therapy self-confidence workshops for people with depression: randomised controlled trial. *Br J Psychiatry*. 2014;204:222-233.
19. Craig AR, Hancock K, Dickson H, Chang E. Long-term psychological outcomes in spinal cord injured persons: results of a controlled trial using cognitive behavior therapy. *Arch Phys Med Rehabil*. 1997;78:33-38.
20. Butler AC, Chapman JE, Forman EM, Beck AT. The empirical status of cognitive-behavioral therapy: a review of meta-analyses. *Clin Psychol Rev*. 2006;26:17-31.
21. Sorensen S, Pinquart M, Duberstein P. How effective are interventions with caregivers? An updated meta-analysis. *Gerontologist*. 2002;42:356-372.

22. Pinquart M, Sorensen S. Helping caregivers of persons with dementia: which interventions work and how large are their effects? *Int Psychogeriatr*. 2006;18:577-595.
23. Brenes GA, Danhauer SC, Lyles MF, Hogan PE, Miller ME. Telephone-Delivered Cognitive Behavioral Therapy and Telephone-Delivered Nondirective Supportive Therapy for Rural Older Adults With Generalized Anxiety Disorder: A Randomized Clinical Trial. *JAMA Psychiatry*. 2015.
24. Kavanagh D, Connolly JM. Mailed treatment to augment primary care for alcohol disorders: a randomised controlled trial. *Drug Alcohol Rev*. 2009;28:73-80.
25. Mains JA, Scogin FR. The effectiveness of self-administered treatments: a practice-friendly review of the research. *J Clin Psychol*. 2003;59:237-246.
26. Chan AW, Tetzlaff JM, Gotzsche PC et al. SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. *BMJ*. 2013;346:e7586.
27. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist*. 1969;9:179-186.
28. Mangione CM, Lee PP, Gutierrez PR et al. Development of the 25-item National Eye Institute Visual Function Questionnaire. *Arch Ophthalmol*. 2001;119:1050-1058.
29. Norman R, Cronin P, Viney R. A pilot discrete choice experiment to explore preferences for EQ-5D-5L health states. *Appl Health Econ Health Policy*. 2013;11:287-298.
30. Martin-Carrasco M, Otermin P, Perez-Camo V et al. EDUCA study: Psychometric properties of the Spanish version of the Zarit Caregiver Burden Scale. *Aging Ment Health*. 2010;14:705-711.
31. Zarit SH, Reever KE, Bach-Peterson J. Relatives of the impaired elderly: correlates of feelings of burden. *Gerontologist*. 1980;20:649-655.

32. Andresen EM, Malmgren JA, Carter WB, Patrick DL. Screening for depression in well older adults: evaluation of a short form of the CES-D (Center for Epidemiologic Studies Depression Scale). *Am J Prev Med.* 1994;10:77-84.
33. Schwarzer R, Jerusalem M. Generalized Self-Efficacy scale. In: Weinman J, Wright S, Johnston M, eds. *Measures in health psychology: A user's portfolio. Causal and control beliefs.* Windsor, UK: NFER-NELSON 1995:35-7.
34. Krupp LB, LaRocca NG, Muir-Nash J, Steinberg AD. The fatigue severity scale. Application to patients with multiple sclerosis and systemic lupus erythematosus. *Arch Neurol.* 1989;46:1121-1123.
35. Herdman M, Gudex C, Lloyd A et al. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res.* 2011;20:1727-1736.

Figure 1. Flowchart of RCT

Table 1. Contents of the 10-week multi-modal treatment - comprising five cognitive behavioural therapy (CBT) modules alternated with five Carers NSW Talk-link group counselling sessions.

Name of the CBT module	CBT Module Inclusions	Talk-link Session
<i>Session One</i> Psychoeducation: AMD education and introduction to CBT	<ul style="list-style-type: none"> - Information regarding AMD. - Misconceptions regarding AMD. - Information regarding the therapeutic intervention, leading to goal setting and expectations. 	<u>Introduction</u> <ul style="list-style-type: none"> • Go through group guidelines and structure to promote safety in the group • Introduction, getting to know one another, sharing individual caring stories • Identifying what members would like to gain from the group most
<i>Session Two</i> Stress response and mood	<ul style="list-style-type: none"> - Education of stress response and mood - Mindfulness, structured relaxation - Relief of anxiety, stress, tension 	<u>Coping with stress</u> <ul style="list-style-type: none"> • Exploring how stress is individually manifested and discussion around individual coping strategies • Encouraging the awareness of individual warning signals that point to the experience of stress • Exploring the concept of mindfulness as a method of stress reduction
<i>Session Three</i> Sleep, general wellbeing and scheduling pleasant events	<ul style="list-style-type: none"> - Basic information and tips regarding nutrition, exercise, socialising and sleep. - Specifically sleep education hygiene, routine. - Encouraging carers to take time to engage in enjoyable/pleasant activities - Designating times to engage in pleasant events 	<u>Self-care</u> <ul style="list-style-type: none"> • Discussing the importance of self-care and individual self-care strategies • Discussion of the importance of our mindset on sleep, relaxation, nutrition and exercise
<i>Session Four</i> Identifying and challenging negative thoughts	<ul style="list-style-type: none"> - Negative thinking/cognitive distortions - Identifying negative thoughts, feelings - Thought recording 	<u>Loss, grief and gain</u> <ul style="list-style-type: none"> • Coping with difficult feelings related to the grief and loss experience and exploring what

	- Thought challenging	<p>carers can ‘gain’ in the process</p> <ul style="list-style-type: none"> • Education about models of grief and loss and discussion of individual experiences • Normalising feelings of anger and sadness • Discussion of individual options for support • Discussion of resilience as positive coping mechanism
<p><i>Session Five</i> Problem solving, resiliency and self-efficacy</p>	<ul style="list-style-type: none"> - Identifying the problem - Identifying and evaluating potential solutions - Problem solving strategy – identifying the best course of action - Interpersonal and communication skills (i.e., assertiveness) - Immediate beliefs and rules - Assumptions around caring - Restructuring core beliefs - Modelling 	<p><u>Communication</u></p> <ul style="list-style-type: none"> • Identifying different communication styles • Communication within the caring role • Discussion of what is going well and what are the difficulties within individual care circumstances • Where to next – resources for further support

Table 2. Validated instruments to determine specific caregiver outcomes collected at baseline; after completion of the 10-week treatment program; and 6 and 12 months post-intervention in family caregivers from both RCT arms.

Scale/ Tool	Approach taken	Outcome evaluated
Caregiver Burden Scale (CBS) ^{12,30}	<ul style="list-style-type: none"> - 22 questions on the impact of the care recipient's disabilities on carer life - Total burden score calculated^{12,31}; higher scores indicate higher levels of caregiver burden 	<ul style="list-style-type: none"> - Assesses how the caregiver perceives the impact of the burden of caregiving (subjective caregiver burden). This is the primary outcome. - Topics covered are: caregiver's health, psychological wellbeing, finances, social life, and relationship between caregiver and care recipient
Centre for Epidemiologic Studies Depression Scale (CESD-10) ³²	-10 items; a score of ≥ 10 out of 30 indicates significant depressive symptoms ³²	- Measures depressive feelings and behaviours experienced in the past week
General Self-Efficacy Scale (GSE) ³³	-10 items; total score ranges between 10 and 40, with a higher score indicating more self-efficacy.	- Assess a general sense of perceived self-efficacy with the aim in mind to predict coping with daily hassles as well as adaptation after experiencing all kinds of stressful life events.
Fatigue Severity Scale ³⁴	- 9 items; a score of ≥ 4 generally means that the fatigue is severe.	- Questionnaire that rates the severity of fatigue symptoms in individuals.
EQ-5D-5L ³⁵	<ul style="list-style-type: none"> - Defines health in terms of 5 dimensions: Mobility; Self-Care; Usual Activities; Pain/Discomfort; and Anxiety/Depression - Used to calculate utility weights and quality adjusted life years (QALYs) during the duration of the trial 	<ul style="list-style-type: none"> - Generic instrument for describing and valuing health will be administered to caregivers - Used for cost-utility analysis